
# avenly Cosmetic

START LOSING

WEIGHT TODAY! We can help you lose 10, 20, 40, or 80 pounds.

## Medical Weight Loss Program \$165 per month reg. \$200

includes Office Visit & Meal Planner, One Month Worth of Medications & Monthly Diet Shot! Our combination of medications promotes weight loss by:



Back or axillary.



#### LIPODISSOLVE

BETTER THAN LIPOSUCTION Gain back your gorgeous look without surgery You can get it all without surgery in as Little as 4 weeks!

# LIPODISSOLVE \$200 PER AMEN' Areas: Abdomen, (Upper or Lower), Love Handles, Low Back, thighs (inner, lateral or posterior), mid

#### SPECIAL PACKAGES

Package A: Lose up to 10 lbs. and up to one inch from waist in 4 weeks. Includes one treatment of Lipodissolve on 3 areas: abdomen (upper or lower), love handles and lower back. One month medical weight loss program. \*Monthly payment of \$55 for 12 months.

\$665

Package B: Lose up to 10-15 lbs and up to 1-2 inches from waist in 4 weeks. Includes 2 Lipodissolve treatments on 3 areas: abdomen (upper or lower), love handles and lower back. One month medical weight loss program. \*Monthly payment of \$97 for 12 months.

NO INTEREST • PAYMENT PLANS FOR 12 MONTHS

**SAVE \$100** 

FROM ANY PACKAGE ABOVE

Expires 6-30-2010

**RESTYLANE®** \$425 per cc

BOTOX® \$160 per area

(Forehead, Glabella, Crows Feet)

\$500 per cc

FINANCING AVAILABLE through "Care Credit" Subject to approval. Results may vary

Saturday Appointments Available.

We Have 2 Locations . Free Parking

STUDIO CITY 11390 Ventura Blvd. . Suite 5 CALL NOW! (818) 361-3788

SAN FERNANDO 501 MacClay Ave. CALL NOW! (818) 361-3788

www.heavenlycosmetic.net



By Dr. Jenn Berman

With celebrities like Brooke Shields, Marie Osmond, Gwyneth Paltrow and Courteney Cox speaking publicly about their experiences with postpartum depression, there is a much greater awareness of the condition than in the past. Yet many mothers are unable to recognize their own symptoms even while they are experiencing it. Family members, spouses, partners and friends often miss the signs, too. The sooner the symptoms of postpartum depression are recognized and treated, the better.

If symptoms last for more than a few weeks or the mother is having thoughts about hurting herself or her baby, she must talk to her OB/GYN immediately and get a referral for a therapist or a psychiatrist who is familiar with postpartum.

## Taking Care of Baby

For the sake of the child and the attachment relationship between parent and infant it is crucial to recognize the problem and get help. Therapy or medication is not a sign of weakness; on the contrary, it is a sign of a good mother who is willing to do whatever it takes to be a good parent.

Studies show depressed moms have the following characteristics:

- Are less active
- Have fewer positive interactions with their children
- Are more likely to poke and prod at their children
- Make less eye contact with their children
- Are less likely to speak "motherese"
- Express more anger
- Mirror their children's vocalizations less frequently
- Are slower to respond to their children
- · Are more disengaged

Studies show children of depressed moms have the following characteristics:

- Explore less
- Play less
- Show fewer positive facial expressions
- Fail to show face and voice preferences
- Make less eye contact
- Don't interact with strangers as well
- Protest more frequently
- Take longer to adjust to new situations
- Are more likely to reverse the mirroring process with their parents resulting in early care-taking behavior



While infants are resilient and have many methods in their repertoire to get their needs met, a mother's depression can have a detrimental effect on her child. Babies are wired to be alert to their mother's emotional states. When a mom is depressed, her face is less expressive, she is less attuned to her child's needs and her energy level is low, all of which have a direct impact on her child. Because of the strong bond between child and mother, this can easily effect a child's cognitive skills, expressive language development and attention span.

Sometimes a mother who is not able to do something for herself, is still able to do that thing for her child. If you, or someone you love, is suffering from postpartum depression, it is crucial for you to get help.

**Dr. Jenn Berman**, a licensed psychotherapist in private practice, is the author of The A to Z Guide to Raising Happy Confident Kids and SuperBaby: The 12 Ways Parents Can Give Their Child a Head Start in the First Three Years. She lives in Los Angeles with her husband and children.

www.DoctorJenn.com

#### Types of **POSTPARTUM DEPRESSION**

#### **Baby Blues**

**Frequency:** 50-85% of new mothers **Occurrence:** 1 day to 2 wks after birth.

**Duration:** 3 wks or less

Symptoms: feeling weepy, moody, irritable, difficulty

concentrating, and/or difficulty sleeping.

#### **Postpartum Depression (PPD)**

**Frequency:** 10-20% of new mothers **Occurrence:** 4 wks to 12 mos after birth

**Duration:** 2 wks to 1 yr.

**Symptoms:** diminished interest in most activities, feelings of being overwhelmed, difficulty concentrating, changes in appetite, guilt, panic attacks, insomnia, weepiness, feelings of worthlessness, excessive worry, fear of being a bad mother, discomfort interacting with the baby, difficulty making even minor decisions and,

occasionally, suicidal thoughts.

#### **Postpartum Psychosis**

Frequency: 0.1- 0.2% of new mothers

**Occurrence:** The two highest risk times for this to occur is in the first 2 wks after childbirth and again at 18-24 mos after delivery.

**Duration:** Sufferers must get immediate medical attention as the condition has a 5% suicide rate and a 4% infanticide rate.

**Symptoms:** Common symptoms include hallucinations, disorientation, confusion, false or bizarre thinking, delusions, fears of the baby being demonic or possessed, and suicidal or homicidal impulses.

